

Factsheet – Natural ponds and artificial surface dams

AUTHORS: E. Lind, S. Filipsson, F. Hedman A. Serra

DATE: 01/03/2021

VERSION: v7



 $\langle 0 \rangle$

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement N°776541.



Natural ponds and artificial surface dams



Unique selling points:

- ✓ Rainwater storage and groundwater replenishment.
- ✓ Reduction of the saline intrusion to the aquifers.

Description of the technology

Natural ponds and artificial surface dams are areas filled with water (such as rainwater, drainage water from **automatic floodgates**, etc.), that are smaller than lakes, and can be used for water storage and infiltration into the aquifers. They are used to manage stormwater runoff, prevent flooding and downstream erosion, and improve water quality in an adjacent river, stream, lake or bay.

They are essentially shallow ponds that are designed to infiltrate stormwater through permeable soils into the groundwater aquifer. Infiltration basins do not release water except by infiltration, evaporation or emergency overflow during flood conditions.

In the NextGen project, natural ponds and artificial surface dams are used to store the rainwater collected and distributed from the **automatic floodgate** ditches. The capacity of these systems is approximately of 8500 m³/ha/year of water storage.

A topographic map where the natural ponds and artificial surface dams will be constructed in the Gotland region can be observed in Figure 1.

Error! No s'ha trobat l'origen de la referència. shows the technology that is being used to find appropriate aquifers where water can be infiltration through the natural ponds and artificial surface.





Flow scheme of the technology



Figure 1. The topographic map to the right shows the soil depth to the rock. As there are so called depressions in the rock and the depression seems to be filled mainly with sand, this area provides high potential for development of a ground water storage system.



Pictures of the technology

Figure 2. The use of electromagnetic Sky-TM measurements from helicopter for finding ground water pounds. The area n^o 2 is the area shown in figure 1 and shows a so-called depression in the rock that most probably is filled mainly with sand and therefore should be a good area for underground storage of water.

Synergetic effects and motivation for the implementation of the technology

✓ Replenishment of groundwater sources

Natural ponds and artificial surface dams are used to store the water directly collected from rainwater in rainy periods and water drained by the **automatic floodgates** to these systems and to gradually infiltrate it to the aquifer, increasing the water availability of the area.

✓ Increase of groundwater quality

Rainwater infiltrated into the aquifer allows reducing the water stress on the local groundwater sources and additionally, in many cases, increases the water quality of the aquifer reducing the salinity of the water.





Natural infiltration ponds and artificial surface dams

✓ Reduction of energy consumption for groundwater pumping

Increasing the water levels of the aquifers through locally infiltration of water allows to avoid long piping distance and will reduce the energy required for long distance pumping of drinking water and could replace investments in desalination which is related to high energy demand.

Requirements of the technology and operating conditions

The following table summarize the requirements of the natural ponds and artificial surface dams to operate in the most appropriated conditions.

Table 1. Required specifications for the natural ponds and artificial surface dams.

Parameter	Units	Min	Max	Reference
Temperature	₽C	n a	20	Below 0 the surface dams will freeze, max 20 (above 20 C, the water has to be chilled by a cooling system).
Rainfall	mm/vear	200	700	(LIVSFS 2017.2) SMHL Vearly precipitation
Water volume	m ³	20 000	95 000	Gotland County Administrative Board. Oral recommendations.
Turbidity	NTU	na	50	SGU, 2013

Key performance indicators

Table 2. KPIs for natural ponds and artificial surface dams.

Parameter	Units	Min	Max	Reference
Percentage of water recovered	%	50	200	NextGen, D1.2
Removal yield of metals	%	-	95	NextGen, D1.2
Removal yield of conductivity	%	-	10	NextGen, D1.2
Removal yield of colour	%	-	70	NextGen, D1.2
Removal yield of COD	%	-	80	NextGen, D1.2
Removal yield of anions	%	-	10	NextGen, D1.2
Removal yield of <i>E.</i> coli	%	-	99	NextGen, D1.2

Links to related topics and similar reference projects

Natural ponds and artificial surface dams	Reference				
NextGen	Case study "Gotland" (NextGen)				



References

Livsmedelsverkets författningssamling (Swedish Food Agency) (2017). Livsmedelsverkets föreskrifter om ändring i Livsmedelsverkets föreskrifter (SLVFS 2001:30) om dricksvatten. LIVSFS 2017:2.

https://www.livsmedelsverket.se/globalassets/om-oss/lagstiftning/dricksvatten---naturlmineralv---kallv/livsfs-2017-2_web.pdf?AspxAutoDetectCookieSupport=1_

Sveriges meteorologiska och hydrologiska institut, SMHI (2020). Klimatdata – kartor. https://www.smhi.se/data/meteorologi/kartor/arsnederbord

Sveriges geologiska undersökning Geological Survey of Sweden, SGU (2013). SGU-rapport 2013:01. *Bedömningsgrunder för grundvatten (Assessment criteria for groundwater)*. ISBN 978-91-7403-193-5. 238p.

http://resource.sgu.se/produkter/sgurapp/s1301-rapport.pdf

Outlook

Case study specific information will be provided, when the results of the other work packages are available:

- Lessons learned from the case study
- Outcome of the assessments
- Legal and regulatory information concerning the whole value chain concerning the technology
- Business opportunities

